

# What you need to know about monkeypox vaccines

The U.S. currently has a limited supply of the vaccine to prevent monkeypox.

The vaccine is a 2-dose series given **before** a risk of exposure **OR** between **4-14 days after** the date of exposure to person who has monkeypox but **before symptoms appear**. Full immunity is reached 2 weeks after the second dose.

The Centers for Disease Control and Prevention and Montana Department of Public Health and Human Services have determined that the following people are most at risk of getting monkeypox and may qualify for a vaccine:

- ★ People who have had a close contact with a person who has monkeypox.
- ★ People with certain risk factors that make them more likely to have been recently exposed to monkeypox.
- ★ People who have recently had multiple or anonymous sexual partners.
- ★ Partners of people who have had multiple or anonymous sexual partners.
- ★ People experiencing homelessness with high-risk behaviors.
- ★ Staff at establishments where sexual activity occurs (e.g. bathhouses, saunas, sex clubs).
- ★ Sex workers.
- ★ People who were diagnosed with gonorrhea or early syphilis within the last 12 months.
- ★ People on HIV pre-exposure prophylaxis.
- ★ People deemed to have high risk based on individual risk assessments.

To learn more about vaccine eligibility, contact RiverStone Health Public Health

406.247.3396 • [phs.info@riverstonehealth.org](mailto:phs.info@riverstonehealth.org)



Scan code to  
request vaccine:



For more info: [cdc.gov/monkeypox](https://cdc.gov/monkeypox)  
or [RiverStoneHealth.org/monkeypox](https://RiverStoneHealth.org/monkeypox)